

# Protecting the Kimberley's Fitzroy River

## Science Statement of Support

Statement from the undersigned coordinating authors and supporting scientists

---

We welcome the Western Australian Government's commitment to create a new protected area in the Fitzroy River Catchment, and to create a management plan to ensure the protection of the National Heritage and other significant cultural and environmental values of the river and provide a basis for sustainable development.

The Fitzroy River is one of the largest principally unregulated rivers remaining in Australia and is recognised as a centre of cultural life and biodiversity in the Kimberley.<sup>1</sup> The river follows a path from the roof of the central Kimberley Plateau, flowing through spectacular and ancient gorges in the King Leopold and Napier Ranges, and coursing between wide floodplains before meeting the ocean at King Sound. The river and catchment support a rich and unique biodiversity of aquatic and terrestrial life with national heritage listed natural and cultural values.<sup>2</sup>

Cultural values arise from unique archaeological records, major rock art assemblages, and associated and continuing Aboriginal cultural practices, and are exceptionally high in the Fitzroy region. Some of the oldest, continuous and most detailed records of Aboriginal occupation, exploitation of both arid and tropical faunal and botanic suites, and persistence during the Last Glacial Maximum come from the Oscar and Napier Ranges<sup>49</sup>. The sites of Carpenter's Gap 1 and Riwi are in the earliest tranche of sites dated by radiocarbon in the continent and have unheralded preservation records of plants, terrestrial and aquatic fauna, organic and inorganic artefacts including some of oldest known edge-ground axes, bone and bifacial Kimberley Points, and marine shell personal ornaments. The rock art of the region is widespread and highly variable and shares strong cultural traditions with the wider Kimberley, having both Pleistocene (pre-10,000 year old style elements) and later styles including variants of Wanjina from the Holocene (during the last-10,000 years). There are also elements of the painted and engraved art bodies, stone arrangements and mythic tracks affiliated with groups of the sandy deserts to the south. Current cultural art estates are actively maintained by Traditional Owners and Indigenous Protected Area Rangers.

The Fitzroy River Declaration by Traditional Owners further speaks to the ongoing multifaceted value of the river to Aboriginal people and outlines aspirations for the future. It is hoped that this statement will also be of value to Traditional Owners in the process of furthering the aspirations expressed in the Declaration.

The Fitzroy River and its estuary provide critical habitat to arguably the world's most important population of Vulnerable (EPBC Act) Freshwater Sawfish (*Pristis pristis*) and supports other elasmobranch species of conservation significance, such as the Freshwater Whipray (*Urogymnus dalyensis*), the Bull Shark (*Carcharhinus leucas*), the Critically Endangered Northern River Shark (*Glyphis garricki*), the Vulnerable Dwarf Sawfish (*Pristis clavata*) and the Winghead Shark (*Eusphyrus blochii*)<sup>3</sup>. Juveniles of the Freshwater Sawfish occur as far upstream in the Fitzroy River as the Margaret River Gorge. Globally, this species has disappeared in >70% of its former range, and intact nurseries, such as the Fitzroy are rare, and are crucial for its persistence globally. In addition, the estuary of the river is a significant nursery for Dwarf Sawfish<sup>4,5,6,7,8</sup>. The aquatic habitat supports

thirty-seven species of fish, many of which are endemic to the Kimberley including an endemic glassfish, archerfish, hardyhead, grunter and gudgeon<sup>9,10,11</sup>.

The lands and waterholes flanking the river are of equally high conservation value. Riparian and floodplain vegetation in the catchment includes 40 species of groundwater-dependent plants, which support riparian bird species such as the nationally Endangered Purple-crowned Fairy Wren and Buff-sided Robin<sup>12,13</sup>. The wetlands and billabongs at the river mouth and along the floodplain are habitat for internationally protected migratory (and non-migratory) birds<sup>13</sup>. In particular, the large floodplain wetlands at Camballin are important waterbird habitat (67 species recorded) and significant breeding grounds for Magpie Geese and Whistling Duck<sup>14</sup>; parts of the floodplain meet criteria for listing as Wetlands of International Importance, under the Ramsar Convention. The river also provides a rich source of food for local Aboriginal and non-Aboriginal people through its abundant fish, freshwater turtle, mussels and freshwater shrimp (cherabin)<sup>15</sup>.

Recent genetic analysis of lizards from across the Kimberley has demonstrated that the Fitzroy region and flanking King Leopold Ranges have a rich fauna, reflecting the mixing of arid zone and tropical elements. In particular, the limestone ranges of the Devonian Reef System<sup>44</sup> flanking the southern margin of the Fitzroy Gorge harbour several endemic species and genetic lineages of lizards, such that this system is a major hotspot of unique evolutionary diversity<sup>45</sup>. Each of these ranges – the Napier, Oscar, Pillara Ranges and scattered outcrops south to Ngumpun Cliffs – are home to unique elements of the biota not found elsewhere<sup>48,46</sup>.

The ecology and hydrology of the river is shaped by highly variable and unpredictable river flows. Waterholes along the course of the river and its tributaries provide vital refugia for wildlife during the dry season months, including fish, turtles, invertebrates, crocodiles and birds. Many waterholes are fed by groundwater during the dry season in a complex interaction between surface and ground water. Surface waters then recharge ground water in the wet season when river flows occur<sup>13, 14,17</sup>. The magnitude of these interactions is likely to vary significantly from year to year, as the volume of and duration of river flow is highly variable. These interactions between surface water and groundwater will also be highly variable along the length of the river, requiring detailed mapping and data collection over at least 5-10 years to characterise given inter-annual variability.

High volume freshwater flows from the river into King Sound are critical for both ecological functioning and the long-term sustainability of the pearling, recreational and commercial fishing and tourism industries. These flows act as a pathway for those species that spend part of their life-cycle in both marine and freshwater environments, such as Barramundi, Freshwater Sawfish and Mulletts. Scientific research has consistently identified the importance of wet season flows from northern Australian rivers in determining the productivity of many species of estuarine fish and crustaceans of economic significance<sup>18, 19,36</sup>.

Significant flood events in the river catchment replenish the vitality and connectivity of the river, floodplains, wetlands, waterholes and estuarine system. More than one third of all fish species in the freshwater reaches of the river need water flows to access the estuary and near shore zone to reproduce, including the ecologically and economically important Barramundi<sup>15, 16</sup>. Tracking of Barramundi in other Northern Australian river systems demonstrates that it is critical to maintain the connectivity between floodplains, the river and estuaries in order to support populations of large-bodied fish<sup>18,19,20</sup>.

With such enviable relatively intact freshwater, riparian and adjacent terrestrial ecosystems that support endemic, endangered and scientifically significant species and communities, the creation of protected areas and a management plan in the Fitzroy River catchment is an opportunity to establish an internationally acclaimed and scientifically endorsed conservation regime. However, the

ecological, cultural and natural values of the river and new protected areas are also tied to the overall condition of the river system which is threatened by agricultural practices<sup>18, 21, 32</sup> and long-running proposals for large-scale irrigation schemes<sup>33,34,41,42,43</sup>, mining and gas extraction<sup>23</sup>.

Large-scale cattle grazing results in soil compaction, degradation and loss of riparian and aquatic vegetation, increased sedimentation, introduction of weeds and potentially significant collective impacts on water quality, hydrology, aquatic ecosystems<sup>20, 24</sup>, as well as on terrestrial vegetation communities<sup>31</sup> and threatened species<sup>32</sup>. The clearing and fragmentation of native vegetation for cropping and associated inputs of pesticides, herbicides, fertilisers and water<sup>23</sup> can lead to severe impacts on freshwater biodiversity.

Native vegetation clearing and water infiltration from irrigation practices can cause groundwater levels to rise and salinisation, as has occurred in the Ord River agricultural area<sup>25, 26</sup> and in the Murray-Darling Basin. Groundwater salinisation has also been linked to the widespread dieback of *Eucalyptus camaldulensis* (River Red Gum) forests of the Murray River floodplain<sup>27</sup>.

Lowering of the water table (through reduction of surface water flows or through groundwater abstraction) can cause pools to become 'disconnected' for longer periods and in extreme cases whole refuge pools can dry-up, threatening the diverse wildlife that relies on groundwater-fed pools. Permanent pools are the principal dry season refugia for freshwater fishes, sawfish, turtles and crocodiles in the river<sup>28</sup>. Vegetation that is groundwater-dependent, such as *Pandanus aquaticus* that provides habitat for the Endangered Purple-crowned Fairy-wren<sup>32</sup> is also likely to suffer from moisture stress<sup>28</sup>.

Stygofauna and troglifauna, which is likely to be diverse in the aquifers of the Fitzroy River catchment, are likewise vulnerable to groundwater extraction<sup>29, 47</sup>. Similarly, lowering of the water table could impact the unique reptiles of the limestone ranges which have persisted here for millions of years at the interface of the desert and Kimberley supported by these complex groundwater systems<sup>13</sup>.

Disruption to natural surface water flows, including damming of the Fitzroy or its tributaries, off-stream storage dams and direct extraction of surface water, may change the frequency and duration with which riparian and floodplain vegetation are inundated by flood waters<sup>30</sup>, and the very morphology of the river system. Water harvesting may have substantial impact on habitat suitability, food-webs and ultimately biodiversity. Damming or re-direction of water flows can interrupt migration and cause isolation (physical and genetic) of populations of aquatic species.

The regular wet-season flooding plays important roles in the recharge of groundwater and therefore persistence of dry-season refuges. In addition, waterbird abundance and composition have already been impacted by such developments in many river and wetland habitats in Australia<sup>28,37</sup>.

Science shows that the most effective way to protect the remarkable natural and cultural values of the river is to: a) create a protected area upstream to eliminate the threat of dams; b) reduce the impacts of landscape-scale threats such as frequent intense fire, weeds and over-grazing that affect water quality; and c) legislate for a management plan that includes a buffer zone along the river that protects important seasonal wetlands; the alluvial aquifers and other groundwater conduits on which dry season refuge pool are reliant, and the floodplains from which the aquifers recharge. The buffer zone must exclude native vegetation clearing, hydraulic fracturing, mining and large water extraction projects<sup>38,39,40</sup>.

The use of good science and modern planning tools should lead to an outcome that matches or exceeds the protection offered in other world-class protected areas in Australia.

## Coordinating Authors

 <p><b>Professor Craig Moritz</b> Australian National University</p>	 <p><b>Professor Kingsley Dixon</b> Curtin University</p>
 <p><b>Professor Peter Veth</b> The University of Western Australia</p>	 <p><b>Professor Jeffery D Connor</b> The University of South Australia</p>
 <p><b>Distinguished Professor Sue O'Connor</b> ARC Laureate Fellow, Centre of Excellence for Australian Biodiversity and Heritage Australian National University</p>	 <p><b>Associate Professor David Morgan</b> Murdoch University</p>
 <p><b>Dr Ryan Vogwill</b> Independent, principal hydrogeologist Previously Associate Professor UWA</p>	 <p><b>Dr Andrew Storey</b> Adjunct Associate Professor The University of Western Australia</p>
 <p><b>Dr Anja Skroblin</b> University of Melbourne</p>	



## Supporting Scientists

<b>Title</b>	<b>Name</b>	<b>Affiliation</b>
Professor	Catherine Lovelock	The University of Queensland
Professor	Hugh Possingham	FNAS, FAA, Dphil (Oxford), The University of Queensland
Professor	John Pandolfi	The University of Queensland
Professor	John Quiggin	School of Economics, The University of Queensland
Professor	John Woinarski	Charles Darwin University
Professor	Keith Christian	Charles Darwin University
Professor	Paul Sunnucks	Monash University
Professor	Peter Davies	Pro Vice-Chancellor (Research), The University of Western Australia
Professor	Pierre Horwitz	Edith Cowan University
Professor	Richard J Hobbs	The University of Western Australia
Professor	Richard Kingsford	University of New South Wales
Professor	Sam Banks	Charles Darwin University
Professor	Tara Martin	University of British Columbia
Professor	Tim Flannery	MSSI, The University of Melbourne
Adjunct Professor	Chris Johansen	The University of Western Australia
Adjunct Professor	William Humphreys	The University of Western Australia
Associate Professor	Jane Balme	The University of Western Australia
Associate Professor	Ben Phillips	The University of Melbourne
Associate Professor	Cynthia Riginos	The University of Queensland
Associate Professor	Jane Elith	The University of Melbourne
Associate Professor	Oscar Venter	University of Northern British Columbia
Emeritus Professor	Stephen Mueke	University of New South Wales
Emeritus Professor	Dale Roberts	The University of Western Australia
Associate Professor	Tim Dempster	University of Melbourne
Dr	A.N. Start	Principle Research Scientist (Retired) DBCA, Research Associate Western Australian Herbarium
Dr	Adrian Gleiss	Murdoch University
Dr	Alexander Watson	Kimberley Program Manager, WWF-Australia
Dr	Alison O'Donnell	The University of Western Australia
Dr	Amelia Wenger	The University of Queensland
Dr	Anne Poelina	Nulungu Research Institute, The University of Notre Dame
Dr	April Reside	The University of Queensland
Dr	Barbara Wueringer	Director and Principle Scientists, Sharks and Rays Australia
Dr	Barry Traill	Director Outback to Oceans Program, Pew Charitable Trusts; Research fellow Charles Darwin University
Dr	Carla Eisemberg	Charles Darwin University
Dr	Cassandra Rowe	James Cook University
Dr	Chandra Salgado Kent	Curtin University

Dr	Christine Adams-Hosking	Global Change Institute, The University of Queensland
Dr	Christine Schlesinger	Charles Darwin University
Dr	Chrystal Mantyka-Pringle	University of Saskatchewan
Dr	David Duncan	The University of Melbourne
Dr	Erica Garcia	TRACK, Charles Darwin University
Dr	Fiona Jean Ede	University of Melbourne
Dr	Jessie Wells	The University of Queensland
Dr	Joe Greet	The University of Melbourne
Dr	Katherine Selwood	University of Melbourne
Dr	Kathy Townsend	University of Sunshine Coast
Dr	Leon A. Barmuta	University of Tasmania
Dr	Matthew Le Feuvre	University of Melbourne
Dr	Michelle L Hall	The University of Melbourne
Dr	Michelle Lim	Adelaide Law School, University of Adelaide
Dr	Neil Pettit	The University of Western Australia
Dr	Nerida Wilson	Western Australian Museum
Dr	Nicole Phillips	The University of Southern Mississippi
Dr	Pauline Grierson	The University of Western Australia
Dr	Peter Kyne	Charles Darwin University
Dr	Reid Tingley	The University of Melbourne
Dr	Renee Gruber	The University of Western Australia
Dr	Rujiporn Thavornkanlapachai	The University of Melbourne
Dr	Sam Nicol	PhD Mathematics / Ecology
Dr	Selina Ward	The University of Queensland, Director (International River Foundation), Councilor (Australian Coral Reef Society)
Dr	Tim Maloney	Griffith University
Dr	Nigel E. Hussey	The University of Windsor
Dr	Yung En Chee	The University of Melbourne
Dr	Gary Scott	University of the Sunshine Coast
Dr	Ross G. Dwyer	The University of Queensland
Dr	Malcolm Lindsay	Kimberley Nature Project
Dr	Mike Christensen	EIANZ, SWCC
Dr	Tegan Davies	The University of Western Australia
Dr	Scott Thompson	EIANZ
Dr	Bill Kruse	Consultant Anthropologist
Dr	Mark Allen	Western Australian Museum
Dr	Glenn Moore	Western Australian Museum
Dr	Simon Clulow	Macquarie University
PhD Candidate	Jordan Iles	The University of Western Australia
PhD Candidate	Glenn Althor	The University of Queensland
PhD Candidate	Jessica Cheok	James Cook University
PhD Candidate	Karissa Lear	Murdoch University
PhD Candidate	Kate Buckley	Charles Darwin University
PhD Candidate	Madeline May Goddard	Charles Darwin University

PhD Candidate	Micha Jackson	The University of Queensland
PhD Candidate	Naomi Indigo	University of Technology Sydney
PhD Candidate	Rosalie Willacy	The University of Queensland
PhD Candidate	Síofra Sealy	Monash University
PhD Candidate	Caroline Mather	The University of Western Australia
	Donna Bagnall	Curtin University
	Dion Wedd	Charles Darwin University
	Andrew Del Marco	Deputy President (WA), EIANZ
	Nigel Jackett	Warden, Broome Bird Observatory
	Steve Wilke	EIANZ
	Simonne Grimes	EIANZ
	George Swann	Kimberley Bird Watching
	Susan Davies	Principal Environmental Consultant, WRM
	Jess Delaney	Principal Environmental Consultant, WRM
	Kim Nguyen	Senior Environmental Consultant, WRM
	Bonita Clarke	Senior Environmental Consultant, WRM
	Chris Hofmeester	Senior Environmental Consultant, WRM
	Emma Thillainath	Environmental Consultant, WRM
	Fintan Angel	Environmental Consultant, WRM
	Adam Harman	Principal Environmental Consultant, WRM



## References

1. Australian Heritage Commission 2011. West Kimberley Place Report: Final Assessment. Retrieved from: <http://www.environment.gov.au/system/files/pages/ed0b4e39-41eb-4cee-84f6-049a932c5d46/files/ahc-final-assessment-full.pdf>
2. Australian Heritage Commission 2011. *ibid*
3. Morgan, D.L., Whitty, J.M., Phillips, N.M., Thorburn, D.C., Chaplin, J.A. and McAuley, R. (2011). North-western Australia as a hotspot for endangered elasmobranchs, with particular reference to sawfishes and the Northern River Shark. *Journal of the Royal Society of Western Australia* 94: 345-358.
4. Morgan, D.L., Somaweera, R., Gleiss, A.C., Beatty, S.J. & Whitty, J.M. (2017). An upstream migration fought with danger: freshwater sawfish fending off sharks and crocodiles. *Ecology* 98: 1465-1467.
5. Phillips, N.M., Chaplin, J.A., Morgan, D.L. & Peverell, S.C. (2011). Population genetic structure and genetic diversity of three critically endangered *Pristis* sawfishes in northern Australian waters. *Marine Biology* 158: 903-915.
6. Phillips, N.M., Chaplin, J.A., Peverell, S.C. & Morgan, D.L. (2017). Contrasting population structures of three *Pristis* sawfishes with different patterns of habitat use. *Marine and Freshwater Research* 68: 452-460.
7. Phillips, N.M., Fearing, A. & Morgan, D.L. (2017). Genetic bottlenecks in *Pristis* sawfishes in northern Australian waters. *Endangered Species Research* 32: 363-372.
8. Whitty, J.M., Keleher, J., Ebner, B.C., Gleiss, A., Simpfendorfer, C.A. & Morgan, D.L. (in press). Habitat use of an intermittent tropical riverine nursery by a critically endangered sawfish. *Endangered Species Research*.
9. Morgan, D.L., Allen, M.G., Bedford, P. and Horstman, M. (2004). Fish fauna of the Fitzroy River in the Kimberley region of Western Australia – including the Bunuba, Gooniyandi, Ngarinyin, Nyikina and Walmajarri Aboriginal names. *Records of the Western Australian Museum* 22: 147-161.
10. Morgan, D.L., Allen, G.R., Pusey, B.J. & Burrows, D.W. (2011). A review of the freshwater fishes of the Kimberley region of Western Australia. *Zootaxa* 2816: 1-64.
11. Morgan, D.L., Unmack, P.J., Beatty, S.J., Ebner, B.C., Allen, M.G., Keleher, J.J., Donaldson, J.A. & Murphy, J. (2014). An overview of the 'freshwater fishes' of Western Australia. *Journal of the Royal Society of Western Australia* 97: 263-278.
12. Kennard, M.J. (ed) (2010). *Identifying high conservation value aquatic ecosystems in northern Australia*. Interim Report for the Department of Environment, Water, Heritage and the Arts and the National Water Commission. Charles Darwin University, Darwin.
13. Pusey, B. and Kath, J. (2015). Environmental Water Management in the Fitzroy River Valley. Information availability, knowledge gaps and research needs. Northern Australia Environmental Resources Hub. Retrieved from: <http://www.waterforfood.wa.gov.au/waterforfood/media/Publications/Fitzroy-River-environmental-water-review-of-data-and-research-needs.pdf>
14. Storey AW, Davies PM and Froend RH (2001). *Fitzroy river system: environmental values*. Report prepared for the Waters and Rivers Commission. Perth, Western Australia.
15. Jackson, S., Finn, M., Woodward, E. and P. Featherston 2011. Indigenous socioeconomic values and river flows, CSIRO Ecosystem Sciences, Darwin.
16. Morgan, D, M G Allen, P Bedford and M Horstman (2002). Inland fish fauna of the Fitzroy River, Western Australia. Report to the Natural Heritage Trust.
17. Harrington, G.A. & Harrington, N.M. (2015). Lower Fitzroy River Groundwater Review. A report prepared by Innovative Groundwater Solutions for Department of Water, May 2015.

18. Robins, J.B., Halliday, I.A, Staunton-Smith, J. Mayer, D.J. and Sellin, N.J. (2005). Freshwater-flow requirements of estuarine fisheries in tropical Australia: a review of the state of knowledge and application of a suggested approach. *Marine and Freshwater Research* 56, 343–360.
19. Pusey BJ, Kennard MJ and Arthington AH. (2004) Freshwater fishes of north-eastern Australia. CSIRO Publishing, Collingwood, Victoria.
20. National Australia Environmental Research Program 2015. Big wet ebbs and flows. Retrieved from: [http://www.nespnorthern.edu.au/wp-content/uploads/2015/11/article\\_nerp\\_crook.pdf](http://www.nespnorthern.edu.au/wp-content/uploads/2015/11/article_nerp_crook.pdf)
21. WA department of Primary Industries and Regional Development, Water for Food – Fitzroy valley groundwater investigations (2017). <http://www.waterforfood.wa.gov.au/Projects/Fitzroy-Valley-Groundwater-Investigation>
22. Harrington, G.A. & Harrington, N.M. (2015). Lower Fitzroy River Groundwater Review. A report prepared by Innovative Groundwater Solutions for Department of Water, May 2015.
23. Department of Mines, Industry Regulation and Safety, WA Government, Petroleum Titles (2017). <http://www.dmp.wa.gov.au/Documents/Petroleum/PD-SBD-GEO-104D.pdf>
24. Pettit NE, Jardine TD, Hamilton SK, Sinnamon V, Valdez D, Davies PM, Douglas MM and Bunn SE. (2012). Seasonal changes in water quality and macrophytes and the impact of cattle on tropical floodplain waterholes. *Marine and Freshwater Research*. 63: 788–800;
25. Smith, A. and Price, A., (2009). Review and assessment of soil salinity in the Ord River Irrigation Area. CSIRO: Water for a Healthy Country National Research Flagship. Rev. A.
26. Ali, R.,J. Byrne and T. Slaven. (2010) Modelling Irrigation and Salinity Management Strategies in the Ord Irrigation Area. *Natural Resources* 1: 34-56. doi:10.4236/nr.2010.11005
27. Cunningham, S. C., J. R. Thomson, R. Mac Nally, J. Read and P. J. Baker. (2011). Groundwater change forecasts widespread forest dieback across an extensive floodplain system. *Freshwater Biology* 56:1494-1508.
28. Warfe, D. M., Pettit, N. E., Davies, P. M., Pusey, B. J., Hamilton, S. K., Kennard, M. J., Townsend, S. A., Bayliss, P., Ward, D. P., Douglas, M. M., Burford, M. A., Finn, M., Bunn, S. E., and Halliday, I. A. (2011). The 'wet-dry' in the wet-dry tropics drives river ecosystem structure and processes in northern Australia. *Freshwater Biology* 56: 2169–2195.
29. Eberhard, S.M. Halse, S.A., Williams, M.R., Scanlon, M.D., Cocking, J. and Barron, H.J. (2009). Exploring the relationship between sampling efficiency and short-range endemism for groundwater fauna in the Pilbara region, Western Australia. *Freshwater Biology* 54: 885– 901
30. Kingsford, R.T. and Porter, J. (2009), Monitoring waterbird populations with aerial surveys – what have we learnt? *Wildlife Research* 36:29-40
31. Legge, S., Kennedy, M., Lloyd, R., Murphy, S., and Fisher, A. (2011). Rapid recovery of mammal fauna in the central Kimberley, northern Australia, following the removal of introduced herbivores. *Austral Ecology* 36, 791-799.
32. Skroblin, A. and Legge, S. (2012). The influence of fine-scale habitat requirements and riparian degradation on the distribution of the purple-crowned fairy-wren (*Malurus coronatus coronatus*) in northern Australia. *Austral Ecology*. 37, 874-884.
33. SMK consultants. (2017) Gogo Station – Surface Irrigation Development Proposal. Referral to Environmental Protection Authority: Supporting Documentation.
34. Adams, V. M., Álvarez-Romero, J. G., Carwardine, J., Cattarino, L., Hermoso, V., Kennard, M. J., Linke, S., Pressey, R. L. and Stoeckl, N. (2014), Planning Across Freshwater and Terrestrial Realms: Cobenefits and Tradeoffs Between Conservation Actions. *Conservation Letters*, 7: 425–440. doi:10.1111/conl.12080
35. Adams, Vanessa M., Setterfield, Samantha A., Douglas, Michael M., Kennard, Mark J. and Ferdinands, Keith B. (2015). Measuring benefits of protected area management: Trends across

realms and research gaps for freshwater systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*,370(1681):1-12.

36. Novak, Peter A., Douglas, Michael M., Garcia, Erica A., Bayliss, Peter and Pusey, Brad J. (2015). A life-history account of *Macrobrachium spinipes* (Schenkel, 1902) (Cherabin) in a large tropical Australian River. *Freshwater Science*,34(2):620-633.
37. King, Alison J., Townsend, Simon A., Douglas, Michael M. and Kennard, Mark J. (2015). Implications of water extraction on the low-flow hydrology and ecology of tropical savannah rivers: an appraisal for northern Australia. *Freshwater Science*,34(2):741-758.
38. Douglas, M., S. Jackson, S. Setterfield, B. Pusey, P. Davies, M. Kennard, D. Burrows, and S. Bunn. 2011. Northern futures: threats and opportunities for freshwater systems. Pages 203-220 in B. Pusey, editor. *Aquatic biodiversity in Northern Australia: patterns, threats and future*. Charles Darwin University Press, Darwin, Northern Territory, Australia.
39. Pittock, J., Finlayson, M., Arthington, A. H., Roux, D., Matthews, J. H., Biggs, H., Harrison, I., Blom, E., Flitcroft, R., Froend, R., Hermoso, V., Junk, W., Kumar, R., Linke, S., Nel, J., Nunes da Cunha, C., Pattnaik, A., Pollard, S., Rast, W., Thieme, M., Turak, E., Turpie, J., van Niekerk, L., Willems, D. and Viers, J. (2015) 'Managing freshwater, river, wetland and estuarine protected areas', in G. L. Worboys, M. Lockwood, A. Kothari, S. Feary and I. Pulsford (eds) *Protected Area Governance and Management*, pp. 569–608, ANU Press, Canberra.
40. R.T. Kingsford, H. Dunn, D. Love, J. Nevill, J. Stein and J. Tait (2005) Protecting Australia's rivers, wetlands and estuaries of high conservation value, Department of Environment and Heritage Australia, Canberra., Product Number PR050823
41. Logan, J. (2017). KIMCO - West Kimberley Land and Water Infrastructure Production and Processing Project
42. Kimberley Development Commission (2014). An Overview of the Pastoral and Agricultural sectors in the Kimberley
43. Regional Development Australia (2016). The Mid and North-Western Australia Investment Portfolio 2016, pp 60-97
44. Playford P.E., Hocking R.M., & Cockbain A.E. (2009) Devonian reef complexes of the Canning Basin, Western Australia. Geological Survey of Western Australia, Perth.
45. Oliver, P.M., Laver, R.J., Martins, F., Pratt, R.C., Hunjan, S., and Moritz, C. 2017. A novel hotspot of vertebrate endemism and an evolutionary refugium in tropical Australia. *Diversity and Distributions* 23, 53-66.
46. Rosauer D. et al. in review. Real-world conservation planning for evolutionary diversity in the Kimberley, Australia, sidesteps uncertain taxonomy. *Conservation Letters* (revision in progress).
47. Wilson and Ponder. 1992. Extraordinary new subterranean isopods (Peracarida: Crusacea) from the Kimberley Region, Western Australia. *Rec. Aust. Mus.* 44(3): 279-298
48. Cameron RAD. 1992. Land snail faunas of the Napier and Oscar Ranges, Western Australia; diversity, distribution and speciation. *Biological Journal of the Linnean Society* 45:271-286.
49. Wood R, Jacobs Z, Vannieuwenhuysse D, Balme J, O'Connor S, Whitau R 2016 Towards an Accurate and Precise Chronology for the Colonization of Australia: The Example of Riwi, Kimberley, Western Australia. *PLoS ONE* 11(9): e0160123.  
<https://doi.org/10.1371/journal.pone.0160123>